VOL. 497 JOURNAL OF CHROMATOGRAPHY DECEMBER 29, 1989

(Biomedical Applications, Vol. 89)

CONTENTS

(Abstracts/Contents List published in Analytical Abstracts, ASCA, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Chromatography Abstracts, Clinical Chemistry Lookout, Current Contents/Physical, Chemical & Earth Sciences, Current Contents/Life Sciences, Deep-Sea Research/Part B: Oceanographic Literature Review, Excerpta Medica, Index Medicus, Mass Spectrometry Bulletin, PASCAL-CNRS, Pharmaceutical Abstracts, Referativnyi Zhurnal, Science Citation Index and Trends in Biotechnology)

Analysis of the major urinary thromboxane metabolites, 2,3-dinorthromboxane B_2 and 11-dehydrothromboxane B_2, by gas chromatography–mass spectrometry and gas chromatography–tandem mass spectrometry
by W.M. Uedelhoven (Munich, F.R.G.), C.O. Meese (Stuttgart, F.R.G.) and P.C. Weber (Munich, F.R.G.) (Received July 17th, 1989) 1

Studies on anabolic steroids. III. Detection and characterization of stanozolol urinary metabolites in humans by gas chromatography–mass spectrometry
by R. Massé, C. Ayotte, H. Bi and R. Dugal (Pointe-Claire, Canada) (Received July 28th, 1989) 17

Quantitative analysis of polyols in human plasma and cerebrospinal fluid
by J. Kusmierz, J.J. DeGeorge, D. Sweeney, C. May and S.I. Rapoport (Bethesda, MD, U.S.A.) (Received August 8th, 1989) 39

Rapid determination of methandrostenolone in equine urine by isotope dilution liquid chromatography–tandem mass spectrometry
by P.O. Edlund, L. Bowers and J. Henion (Ithaca, NY, U.S.A.) and T.R. Covey (Thornhill, Canada) (Received August 30th, 1989) 49

High-performance liquid chromatographic determination of metabolic products for fermentation control of mammalian cell culture: analysis of carbohydrates, organic acids and orthophosphate using refractive index and ultraviolet detectors
by F. Weigang, M. Reiter, A. Jungbauer and H. Katinger (Vienna, Austria) (Received August 24th, 1989) 59

Isocratic high-performance liquid chromatographic determination of the concentration and specific radioactivity of phosphoenolpyruvate and uridine diphosphateglucose in tissue extracts
by A. Giaccari and L. Rossetti (San Antonio, TX, U.S.A.) (Received August 21st, 1989) 69

Study on the analogy of gel permeation chromatography and cuprophan dialysis for the isolation of uremic 'middle molecules' using model compounds
by A.C. Schoots and C.A.M.G. Cramers (Eindhoven, The Netherlands) (Received September 1st, 1989) 79

(Continued overleaf)

Contents (continued)

Potential of the sulfobetaine detergent Zwittergent 3-12 as a desorbing agent in biospecific and bioselective affinity chromatography
by K.H. Nollstadt, Y.D. Karkhanis, M.J. Gnozzio, M.St.J. Crane, A.M. Gurnett, D.M. Schmatz and M.J. Turner (Rahway, NJ, U.S.A.) (Received August 11th, 1989) 87

Determination of urinary orotate excretion by high-performance liquid chromatography
by V. Ferrari, G. Giordano, A.T. Cracco, N. Dussini, L. Chiandetti and F. Zacchello (Padova, Italy) (Received September 4th, 1989) 101

Detection and separation of intracellular 1-β-D-arabinofuranosylcytosine-5-triphosphate by ion-pair high-performance liquid chromatography
by E. Schleyer (Münster, F.R.G.), G. Ehninger (Tübingen, F.R.G.), M. Zühlsdorf (Münster, F.R.G.), B. Proksch (Tübingen, F.R.G.) and W. Hiddemann (Münster, F.R.G.) (Received August 9th, 1989) 109

Determination of 2-*n*-octadecylindole-5-carboxylic acid in human plasma by high-performance liquid chromatography
by G.D. Jones, C.D. Evans, V. Facchini and O.R.W. Lewellen (Dagenham, U.K.) (Received August 2nd, 1989) 121

Measurement of K vitamins in animal tissues by high-performance liquid chromatography with fluorimetric detection
by K. Hirauchi, T. Sakano, S. Notsumoto, T. Nagaoka, A. Morimoto, K. Fujimoto, S. Masuda and Y. Suzuki (Osaka, Japan) (Received July 20th, 1989) 131

New and versatile method for the determination of faecal bile acids by thin-layer chromatography with direct scanning fluorimetry
by M. Kindel, H. Ludwig-Koehn and B. Lembcke (Göttingen, F.R.G.) (Received August 15th, 1989) 139

Determination of rimantadine and its hydroxylated metabolites in human plasma and urine
by F.A. Rubio, N. Choma and E.K. Fukuda (Nutley, NJ, U.S.A.) (Received September 1st, 1989) 147

Gas chromatography of simple phenols in biological fluids
by M. Balíková and J. Kohlíček (Prague, Czechoslovakia) (Received August 11th, 1989) 159

Liquid chromatographic–atmospheric pressure ionization mass spectrometric analysis of toremifene metabolites in human urine
by N. Watanabe, T. Irie, M. Koyama and T. Tominaga (Tokyo, Japan) (Received August 15th, 1989) 169

Rapid high-performance liquid chromatographic method for the measurement of the enantiomers of metoprolol in serum using a chiral stationary phase
by D.R. Rutledge and C. Garrick (Detroit, MI, U.S.A.) (Received August 11th, 1989) 181

Determination of the enantiomers of verapamil and norverapamil in serum using coupled achiral–chiral high-performance liquid chromatography
by Y.-A. Chu and I.W. Wainer (Memphis, TN, U.S.A.) (Received August 22nd, 1989) 191

Analysis of benzodiazepines and tricyclic antidepressants in serum using a common solid-phase clean-up and a common mobile phase
by M. Mazhar and S.R. Binder (Hercules, CA, U.S.A.) (Received August 28th, 1989) 201

Analysis of albuterol in human plasma based on immunoaffinity chromatographic clean-up combined with high-performance liquid chromatography with fluorimetric detection
by H. Ong, A. Adam, S. Perreault, S. Marleau, M. Bellemare and P. du Souich (Montreal, Canada) and N. Beaulieu (Ottawa, Canada) (Received August 25th, 1989) ... 213

Determination of oxiracetam in plasma and urine by column-switching high-performance liquid chromatography
by J.B. Lecaillon, C. Souppart, F. Le Duigou and J.P. Dubois (Rueil-Malmaison, France) (Received August 28th, 1989) 223

Notes

Das Erythrocruorin des Regenwurms (*Lumbricus terrestris*) als Eichsubstanz in der Gelchromatographie
von W.K.R. Barnikol, O. Burkhard und H. Pötzschke (Mainz, B.R.D.) (Eingegangen am 16. Augustus 1989) 231

High-performance liquid chromatographic determination of branched-chain α-keto acids in serum using immobilized leucine dehydrogenase as post-column reactor
by N. Kiba, M. Muto and M. Furusawa (Kofu, Japan) (Received July 28th, 1989) . 236

A new rapid method for phospholipid separation by high-performance liquid chromatography with light-scattering detection
by L. Breton, B. Serkiz, J.-P. Volland and J. Lepagnol (Suresnes, France) (Received August 11th, 1989) 243

Use of anion-exchange resin in F^- form in sample processing for determination of carnitine
by A. Sandor, J. Cseko and I. Alkonyi (Pecs, Hungary) (Received August 31st, 1989) 250

Isolierung eines Fibronektin-Rezeptors von *Streptococcus pyogenes* durch Affinitätschromatographie und anschliessender Hochleistungflüssigkeitschromatographie
von R. Sting, M. Schmitt und H. Blobel (Giessen, B.R.D.) (Eingegangen am 5. September 1989) 258

Assay for lysine-ketoglutarate reductase by reversed-phase high-performance liquid chromatography
by A.T. Davis (Grand Rapids, MI, U.S.A.) (Received September 4th, 1989) 263

(Continued overleaf)

Contents (continued)

Purification of microbial uricase
by V. Adámek, B. Králová, M. Šüchová, O. Valentová and K. Demnerová (Prague, Czechoslovakia) (Received August 22nd, 1989) 268

Determination of ascorbic acid and dehydroascorbic acid in plasma and cerebrospinal fluid by liquid chromatography with electrochemical detection
by E. Nagy and I. Degrell (Debrecen, Hungary) (Received July 2nd, 1989) 276

Picogram level determination of medetomidine in dog serum by capillary gas chromatography with negative ion chemical ionization mass spectrometry
by L. Vuorilehto, J.S. Salonen and M. Anttila (Turku, Finland) (Received August 14th, 1989) 282

Quantitative gas chromatographic–mass spectrometric analysis of acrivastine and a metabolite in human plasma
by S.Y. Chang, F.R. Nelson, J.W.A. Findlay and L.C.E. Taylor (Research Triangle Park, NC, U.S.A.) (Received September 1st, 1989) 288

Assay of a calcium antagonist (PU 122) in plasma by gas chromatography
by A. Marzo, E. Treffner, G. Meroni and M. Ripamonti (Tribiano, Italy), R. Pelloso (Milan, Italy) and C. Lucarelli (Rome, Italy) (Received July 31st, 1989) 296

High-performance liquid chromatographic assay for cytosine arabinoside and uracil arabinoside in cerebrospinal fluid and plasma
by A. Riccardi, T. Servidei, A. Lasorella and R. Riccardi (Rome, Italy) (Received July 7th, 1989) 302

Measurement of low (sub-therapeutic) phenobarbitone levels in plasma by high-performance liquid chromatography: application to patient compliance studies
by S. Peaker, A.C. Mehta, S. Kumar and M. Feely (Leeds, U.K.) (Received July 19th, 1989) 308

Measurement of underivatised metoprolol enantiomers in human plasma by high-performance liquid chromatography with a chiral stationary phase
by M.S. Ching, M.S. Lennard, A. Gregory and G.T. Tucker (Sheffield, U.K.) (Received July 25th, 1989) 313

High-performance liquid chromatographic study of GDP-mannose and GDP-fucose metabolism
by A. Martin, D. Ruggiero-Lopez, P. Broquet, M. Richard and P. Louisot (Oullins, France) (Received August 31st, 1989) 319

Determination of morphine in plasma by high-performance liquid chromatography with fluorescence detection
by J.J. Schneider and P.J. Ravenscroft (Brisbane, Australia) (Received August 24th, 1989) 326

High-performance liquid chromatographic determination of ibuprofen in rat and human plasma
by J.H. Satterwhite and F.D. Boudinot (Athens, GA, U.S.A.) (Received August 24th, 1989) 330

Mitomycin C determination using loop-column extraction: a rapid and sensitive high-performance liquid chromatographic assay for pharmacokinetic studies with Spherex starch particles
by M.J. Czejka, W. Jäger and J. Schüller (Vienna, Austria) (Received June 14th, 1989) 336

Determination of saterinone in plasma by high-performance liquid chromatography
by M. Rudolph (Hamburg, F.R.G.) (Received July 27th, 1989) 342

Rapid analysis of ceftetrame in human plasma using sorbent extraction and high-performance liquid chromatography
by C.M. Hicks and M.L. Powell (Raritan, NJ, U.S.A.) (Received August 2nd, 1989) 349

Determination of flumequine in fish by high-performance liquid chromatography and fluorescence detection
by O.B. Samuelsen (Bergen, Norway) (Received August 11th, 1989) 355

Letters to the Editor

Production of tyrosine isomers in mice by therapeutic doses of ^{60}Co irradiation
by R.C. Brown, C.E. Evans and J.A. Hoskins (Carshalton, U.K.) and J. Peacock (Belmont, U.K.) (Received July 24th, 1989) 360

Ultra-rapid assay of brain γ-aminobutyric acid by liquid chromatography with electrochemical detection
by S. Murai, H. Saito, H. Nagahama, H. Miyate, Y. Masuda and T. Itoh (Morioka, Japan) (Received August 11th, 1989) 363

Improved method for the determination of furosemide in plasma by high-performance liquid chromatography
by W. Radeck and M. Heller (Frankfurt/Main, F.R.G.) (Received August 18th, 1989) 367

Related articles published in the Journal of Chromatography, Vols. 482 and 483 371

Author Index 373

Subject Index 377

In articles with more than one author, the name of the author to whom correspondence should be addressed is indicated in the article heading by a 6-pointed asterisk (*)